

PTSD LCM Clinician Ratings (RETROSPECTIVE)

Patient Name _____ Years 19__ - 19__

PLEASE PRINT

Other ()	Monthly Medication Dose (if available)	
Other ()		
Other ()		
Thyroid (T3 or T4)		
Benzodiazepine ()		
Neuroleptic ()		
MAOI ()		
Antidepressant II ()		
Antidepressant I ()		
Therapy ()		

Intrusion/ Arousal		
SEVERE	Essentially Incapacitating	SEVERE
MODERATE	Notable Intrusion or Arousal	MODERATE
MILD	Minimal Intrusion or Arousal	MILD
Months of Year → 1 2 3 4 5 6 7 8 9 10 11 12 1 2 3 4 5 6 7 8 9 10 11 12 1 2 3 4 5 6 7 8 9 10 11 12 1 2 3 4 5 6 7 8 9 10 11 12 Baseline		
MILD	Minimal Avoidance or Numbing	MILD
MODERATE	Notable Avoidance or Numbing	MODERATE
SEVERE	Essentially Incapacitating	SEVERE

Avoidance/ Numbing

TRACK COMORBID SYMPTOMS HERE →

- 0= none
- 1= mild
- 2= low moderate
- 3= high moderate
- 4= severe

Severity of Substance Abuse	
Functional Impairment	
Depression	

Impact (-4 to +4)	19__	19__	19__	19__	19__
Life Events					
(Date)					